



(12) **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention
of the grant of the patent:
01.02.2006 Bulletin 2006/05

(51) Int Cl.:
C12N 15/11 (2006.01) A61K 48/00 (2006.01)
A61K 31/70 (2006.01) C12Q 1/68 (2006.01)

(21) Application number: **99937069.5**

(86) International application number:
PCT/JP1999/004399

(22) Date of filing: **13.08.1999**

(87) International publication number:
WO 2000/009684 (24.02.2000 Gazette 2000/08)

(54) **NUCLEIC ACID CAPABLE OF BINDING SPECIFICALLY TO Ras TARGET PROTEIN**

NUKLEINSÄURE, DIE SPEZIFISCH AN EIN RAS ZIELPROTEIN BINDEN KANN

ACIDE NUCLEIQUE CAPABLE DE SE LIER SPECIFIQUEMENT A UNE PROTEINE CIBLE RAS

(84) Designated Contracting States:
CH DE FR GB LI

(56) References cited:
WO-A-92/14843 WO-A-94/29727
WO-A-97/34146

(30) Priority: **14.08.1998 JP 24259698**
24.11.1998 JP 33328498

(43) Date of publication of application:
09.08.2000 Bulletin 2000/32

(73) Proprietor: **Japan Science and Technology**
Agency
Kawaguchi-shi
Saitama (JP)

(72) Inventors:
• **YOKOYAMA, Shigeyuki**
Tokyo 113-0023 (JP)
• **HIRAO, Ichiro**
Asaka-shi,
Saitama-ken 351-0036 (JP)
• **SAKAMOTO, Kensaku**
Tokyo 120-0044 (JP)

(74) Representative: **Cresswell, Thomas Anthony**
J.A. KEMP & CO.
14 South Square
Gray's Inn
London WC1R 5JJ (GB)

- **DATABASE WPI Section Ch, Week 199519**
Derwent Publications Ltd., London, GB; Class
B04, AN 1995-129001 XP002246369 ZENECA
PHARM. CANCER RES. DEPT.; ELLSTON J.M.;
HUDSON K.: "Screening for inhibitors or
ras-protein partner interactions - using a
ras-binding domain coated on scintillation plates
and tritium-labelled GTP" & RESEARCH
DISCLOSURE, vol. 371, no. 034, 10 March 1995
(1995-03-10), Emsworth, GB
- **MICHIKO KIMOTO ET AL.: 'RNA aptamers that**
specifically bind to the Ras-binding domain of
Raf-1' FEBS LETTERS, vol. 441, no. 2, December
1998, pages 322 - 326, XP002926506
- **WALTER KOLCH ET AL.: 'Inhibition of Raf-1**
signaling by a monoclonal antibody, which
interferes with Raf-1 activation and with Mek
substrate binding' ONCOGENE, vol. 13, no. 6,
1996, pages 1305 - 1314, XP002926507
- **ANDREW D. ELLINGTON ET AL.: 'In vitro**
selection of RNA molecules that bind specific
ligands' NATURE, vol. 346, no. 6287, 1990, pages
818 - 822, XP002926508
- **CRAIG TUERK ET AL.: 'Systematic evolution of**
ligands by exponential enrichment: RNA ligands
to bacteriophage T4 DNA polymerase' SCIENCE,
vol. 249, no. 4968, 1990, pages 505 - 510,
XP002926509

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

DescriptionTechnical Field

5 **[0001]** The present invention relates to novel nucleic acids (aptamer) which are specifically bound to target proteins of Ras. More preferably, the present invention relates to novel RNA aptamers which are specifically bound to Raf-1. Further, the present invention relates to control of signal transduction that induces proliferation or differentiation of cells using the nucleic acids of the present invention, and to a pharmaceutical composition using the same.

10 Background of the Invention

[0002] Ras is a guanine nucleotide binding protein, and participates in signal transduction of cells. When a receptor of cells is activated, "GDP binding Ras" in cells becomes "GTP binding Ras".

15 **[0003]** This "GTP binding Ras" is bound to "target proteins of Ras" such as Raf-1, B-Raf, RGL, Ral GDS, MEKK, P13K and the like. These "target proteins of Ras" have a Ras binding domain (RBD) to which the GTP binding Ras can be bound, and the GTP binding Ras is bound to this domain of these "target proteins of Ras" to transmit necessary signals into cells.

[0004] Ras is a key protein of intracellular signal transduction, and the "target proteins of Ras", such as Raf-1, are a center of the intracellular signal transduction system in which signals from Ras are transmitted according to the types.

20 **[0005]** Accordingly, a substance capable of specifically blocking the binding domain with the GTP binding Ras in the "target proteins of Ras", if any, can specifically inhibit an intracellular signal transduction system by Ras, and it is useful to treat or prevent various diseases triggered by the signal transduction. For example, with respect to tumor cells, proliferation or differentiation of tumor cells can be inhibited by specifically controlling the signal transduction that induces proliferation or differentiation with the "target cells of Ras" to treat cancers or inhibit metastasis.

25 **[0006]** By the way, Ras-1, one of the "target cells of Ras" is a serine/threonine protein kinase present in a cytoplasm, and the activity is induced by interaction with the GTP binding Ras. The activated Raf-1 phosphorylates MEK (MAPK/ERK kinase), and then MEK phosphorylates ERK to transmit signals into a nucleus (Daum, G., et al., (1994) Trends Biochem. Sci. 19, 474 - 480; Avruch, J., et al., (1994) Trends Biochem. Sci. 19, 279 - 283).

30 **[0007]** In order to elucidate such an intracellular signal transduction system of Raf-1, a method of selectively inhibiting the function of Ras or Raf-1 has been utilized (deVries-Smits, A. M., et al., (1992) Nature 357, 602 - 604). These studies include inhibition of the Ras function with a Raf-1 mutant free from a kinase activity (Kolch, W., et al., (1991) Nature 349, 426 - 428), inhibition of a Raf-1 kinase with an antibody bound to a kinase domain of Raf-1 (Kolch, W., et al., (1996) Oncogene 13, 1305 - 1314) and the like.

35 **[0008]** However, these inhibitors do not specifically inhibit a specific part of a signal transduction system with Ras or Raf-1, but inhibit many functions such as a function of binding to Ras, a kinase function and the like simultaneously and diversely. Accordingly, a signal transduction system to be inhibited cannot be specified. Thus, individual specific mechanisms of a signal transduction system could not be clarified satisfactorily.

[0009] Consequently, the development of a molecular species capable of specifically inhibiting the binding of Ras to Raf-1 has become important for clarifying the role of the signal transduction system.

40 **[0010]** At present, a downstream signaling pathway of Ras has not been completely clarified. When such a molecular species is developed, it is possible to elucidate the signaling pathway in which Ras participates using a molecular species capable of specifically inhibiting some specific routes and clarify the signaling pathway with target proteins of Ras in detail. In addition, it is possible to control the intracellular signal transduction. Consequently, various diseases in which the intracellular signal transduction participates, such as tumors and the like, can be treated and prevented.

45 **[0011]** Meanwhile, the structural analysis of the "target proteins of Ras" in the intracellular signaling pathway in which Ras participates has been conducted. It has been known that the Ras binding domain (RBD) of Raf-1 is located from 51 to 131 residues in the N-terminus of Raf-1 (Vojtek, A. B., et al., (1993) Cell 74, 205 - 214; Chuang, E., et al., (1994) Mol. Cell. Biol. 14, 5318 - 5325).

50 **[0012]** Further, nucleic acid molecular species (aptamers), such as an RNA, a DNA and the like, having a high affinity for a certain target, such as proteins, have been isolated by "in vitro selection" methods (Ellington, A. D. et al., (1990) Nature 346, 818 - 822; Tuerk, C. et al., (1990) Science 249, 505 - 510) (Bock, L. C., et al., (1992) Nature 355, 564 - 566; Qiu Qiu, Y. L., et al., (1994) Nucleic Acids Res. 22, 5229 - 5234; Gal. S. W., et al., (1998) Eur. J. Biochem. 252, 553 - 562; Bell, S. D., et al., (1998) J. Biol. Chem. 273, 14309 - 14314). Therefore, there is a possibility that an RNA specifically bound to Raf-1 is obtained by applying this method to Raf-1, while the interaction with an RNA is unknown.

55 **[0013]** WO 92/14843 discloses aptamers specific for biomolecules and method of making. This document discloses a method for generating aptamers that bind to particular molecules.

[0014] WO 94/29727 discloses a method of evaluating an anti-proliferative compound by contacting the compound with Raf and determining its ability to bind to Ras.

Disclosure of the Invention

[0015] The present invention is to provide nucleic acid molecular species which can specifically inhibit the binding to "GTP binding Ras" by being specifically bound to a Ras binding domain (RBD) of target proteins of Ras such as Raf-1.

[0016] In order to clarify the signaling pathway in which the target proteins of Ras participate and the physiological activity provided by the inhibition of the signal transduction, the development of substances which specifically inhibit the binding to Ras having an important role in cells in particular along with Raf-1 by being specifically bound to the target proteins of Ras, in more detail, Raf-1 and which have a strong activity have been in demand. The present inventors have found that a nucleic acid molecular species specifically bound to the Ras binding domain of the "target proteins of Ras" can be obtained using the *in vitro* selection method. For example, it has been possible to obtain a novel RNA aptamer targeting the Ras binding domain (RBD) of Raf-1, one of the "target proteins of Ras" by this method and to determine the RNA sequence thereof. This RNA aptamer can specifically inhibit the binding between Ras and Raf-1.

[0017] Accordingly, the present invention is to provide novel nucleic acid molecular species bound to a "target protein of Ras", especially, its Ras binding domain (RBD), an agent for controlling a signal transduction system using the nucleic acid molecular species, a method of controlling the same, and a pharmaceutical composition containing the same.

[0018] More specifically, the present invention is to provide RNA aptamers that bound to a "target protein of Ras", especially its RBD, an agent for controlling a signal transduction system using this RNA aptamer, a method of controlling the same, and a pharmaceutical composition containing the same.

[0019] Accordingly, the invention provides an RNA capable of being specifically bound to a Ras binding domain of Raf-1, wherein the RNA contains at least one sequence selected from sequence Nos. 1 to 8 or 25 to 28 of the Sequence Listing or a sequence of 81 to 99 bases obtainable by deletion of a base or bases from the 3-terminus of sequence No. 25.

Brief Description of the Drawings

[0020]

Fig. 1 shows a sequence of an initial RNA pool in *in vitro* selection and sequences of PCR primers.

Fig. 2 shows 24 sequences of RNAs obtained from the RNA pool in the 21st round. In the drawing, a random sequence moiety of approximately 60 bases is shown. The overall sequence includes the sequences of the 5'- and 3'- termini defined in Fig. 1. Sequences which are the same as that of clone 1 are termed "group 1". The other sequences are termed "group 2". Incidentally, in note (a), the number of clones of ligands isolated respectively is shown in parentheses. In note (b), clone 21.08 indicates that each sequence defined has two mutations. In note (c), percentage of the binding of the RNA ligand to GST-RBD is based on a value measured by the nitrocellulose filter binding assay.

Fig. 3 shows the binding of the RNA ligand to the GST-RBD protein. Percentage of the binding of the RNA ligand to GST-RBD is based on a value measured by the nitrocellulose filter binding assay. In Fig. 3, a closed circle indicates the use of an RNA of sequence No. 1, a closed square the use of an RNA of sequence No. 7, and a closed triangle the use of an RNA of sequence No. 11.

Fig. 4 is a photo that replaces a drawing, showing inhibition of interaction between Ras and GST-RBD by the RNA aptamers. An amount of the Ras protein bound to GST-RBD as measured by immunoblotting with anti-Ras antibody RAS004 is shown on an upper column. An total amount of GST-RBD measured by staining with Coomassie Blue is shown on a lower column. GDP binding (D) or GTP γ S binding (T) Ras (2 pmols) and GST-RBD (25 pmols) were incubated in the presence of an RNA in various amounts. A used 21.01 ligand in Fig. 2, B 21.07 ligand in Fig. 2, C 21.11 ligand in Fig. 2, and D 21.12 ligand in Fig. 2.

Best Mode for Carrying Out the Invention

[0021] The "target proteins of Ras" refers to Ras proteins which participate in the cell signal transduction, preferably a group of proteins forming an intracellular signal transduction system by interacting with GTP binding Ras proteins. Examples of the "target proteins of Ras" included Raf-1, B-Raf, RGL, Ral, GDS, MEKK, P13K and the like. The "target proteins of Ras" of the present invention are preferably Raf-1 and the like.

[0022] The present invention has clarified that nucleic acid molecular species capable of being specifically bound to the foregoing "target proteins of Ras" exist. Accordingly, the nucleic acid specifically bound to the "target proteins of Ras" may be an RNA or a DNA. The RNA or the DNA is not particularly limited so long as it is specifically bound to the "target proteins of Ras". Further, the nucleic acid may be specifically bound to only one "target protein of Ras" or to two or more "target proteins of Ras".

[0023] The size of bases of the nucleic acid molecular species of the present invention is not particularly limited so long as it is sufficient to allow the specific binding to the "target proteins of Ras". It is between 20 and 300 bases,

preferably between 20 and 150 bases, more preferably between 30 and 150 bases, further preferably between 50 and 150 bases. In case the binding specificity is stressed, the longer size is preferable. However, in view of the ease of the procurement in the synthesis method or the like, the shorter size is preferable.

[0024] As the nucleic acid molecular species of the present invention, an RNA containing any one of base sequences, sequence Nos. 1 to 28, preferably sequence Nos. 1 to 8 or sequence Nos. 25 to 28 of Sequence Listing is mentioned.

[0025] The RNAs of the present invention shown in the sequence numbers have an ability of binding to the "target proteins of Ras". More specifically, the RNAs are characterized in that they are specifically bound to the Ras binding domain (RBD) of Raf-1, and the nucleic acid molecular species of the present invention is not limited to the base sequences shown in the foregoing sequence Nos. 1 to 28.

[0026] The nucleic acid molecular species of the present invention can be one containing these base sequences in the whole or a part of the molecule. For example, a base sequence or another molecular species may further be added to the nucleic acid molecular species having the base sequences shown in sequence Nos. 1 to 28 of Sequence Listing unless the ability of binding to the "target proteins of Ras" is inhibited.

[0027] These RNAs of the present invention can also be reversely transcribed, as required, into DNAs having complementary base sequences to the RNAs. Accordingly, the present invention relates to nucleic acid molecular species such as RNAs, DNAs and the like, containing any one of base sequences of sequence Nos. 1 to 28 of Sequence Listing or a base sequence in which at least one base thereof is deleted and substituted with another base and/or at least one base is added.

[0028] The "aptamer" refers to a nucleic acid molecular species capable of being bound to a specific domain of a protein, and the nucleic acid may be an RNA or a DNA. An aptamer made of an RNA is called an "RNA aptamer". Accordingly, in the present invention, the nucleic acid molecular species in the present invention is also termed an "aptamer". When the nucleic acid is an RNA, it is called an "RNA aptamer".

[0029] Incidentally, RNAs comprising approximately 60 bases shown in sequence Nos. 29 to 52 of Sequence Listing indicate base sequences comprising approximately 60 bases in a central portion of RNAs shown in sequence Nos. 1 to 24. Further, RNAs comprising approximately 45 bases shown in sequence Nos. 53 and 54 of Sequence Listing indicate base sequences comprising approximately 45 bases in a central portion of RNAs shown in sequence Nos. 25 to 28. Moreover, sequence Nos. 55 to 60 of Sequence Listing show base sequences of primers used in specific examples.

[0030] The nucleic acid (aptamer) of the present invention can be produced by various methods. When the base sequence of the aptamer is known, it can be synthesized.

[0031] When the base sequence of the aptamer of the present invention is unknown, the aptamer can be produced through selection by the known "in vitro selection" method (Ellington, A. D. et al., (1990) Nature 346, 818 - 822; Tuerk, C. et al., (1990) Science 249, 505 - 510). The "in vitro selection" method is described.

[0032] First, RNAs containing a random base sequence of 20 to 300 bases, preferably 30 to 100 bases, more preferably 30 to 70 bases are produced. These RNAs are produced by transcription from synthetic DNAs containing a random sequence.

[0033] A base sequence which is to be a primer in the PCR method is added to the 5'-terminus and the 3'-terminus of the DNAs. In this case, the primer is not particularly limited. A primer having a sequence of cleavage with a restriction endonuclease so as to be able to cleave this primer portion later is preferable. A size of the primer portion is not particularly limited. It is approximately between 20 and 50 bases, preferably between 20 and 30 bases. Further, the primer at the 5'-terminus may be designed such that a promoter sequence of a T7 RNA polymerase is added thereto to enable the transcription reaction from DNA to RNA.

[0034] In this manner, the RNA group (RNA pool) having the base sequences as the primer at both termini and the random base sequence in the center is produced by transcription of the DNA.

[0035] Subsequently, the RNA in this RNA pool and the "target protein of Ras", for example, Raf-1 or a peptide comprising its binding domain are contacted to separate the RNA bound to the "target protein of Ras". The selected RNA is converted to a cDNA through reverse transcription, and it is amplified by PCR using the primers. The DNA amplified is transcribed into an RNA, and this is returned to the RNA pool.

[0036] One cycle, termed a "round", comprises binding with the "target protein" of Ras in the RNA pool, the separation of the bound RNA, reverse transcription, amplification by PCR and transcription of the DNA. That is, one round means that the foregoing round is conducted once.

[0037] When the foregoing round using the RPTA pool is conducted, the amount of the RNA bound to the "target protein of Ras" in the RNA pool is increased, and further the amount of the RNA having the specific binding base sequence is increased, so that the RNA to be specifically bound can be selected by repeating the round.

[0038] Such a round is conducted 5 to 50 times, preferably 5 to 30 times.

[0039] The RNA sequences selected by the "in vitro selection" method as described above are determined by a usual method, and this RNA can also be converted to a cDNA through reverse transcription by a usual method. Further, the primer regions can be cleaved as required. In this manner, the aptamers of the present invention can be obtained.

[0040] The "in vitro selection" method is described in more detail by using Raf-1 as the "target protein of Ras".

[0041] The present inventors have prepared a pool of RNAs having a random sequence of approximately 60 bases to select RNAs bound to the Ras binding domain (RBD) of Raf-1. And, the base sequences shown in Fig. 1 were bound to the 3'-terminus and the 5'-terminus of these RNAs. It was presumed that approximately 8×10^{13} sequences of RNAs are present in this RNA pool.

5 [0042] Before conducting the selection of the aptamer, the effect of the salt concentration relative to the binding between the RNA pool and Raf-1-RBD was examined. At a low salt concentration, the RNA was non-specifically bound to Raf-1-RBD. However, it was found that the non-specific binding is suppressed by increasing the salt concentration (up to approximately 150 mM). Thus, the present inventors used a phosphate buffer (hereinafter referred to as a "binding buffer") containing 137 mM sodium chloride as a buffer for the selection.

10 [0043] The selection from the 1st to 13th rounds was conducted by the binding between a fusion protein of glutathione S-transferase (GST) and a peptide (RBD) of 51 to 131 of Raf-1 (hereinafter referred to as "GST-RBD") and RNAs using a glutathione-Sepharose 4B matrix. In the 13th round, the binding ability (binding ratio) to the RNA pool was slightly changed from 0.16% at the initial stage to 0.36%.

15 [0044] Successively, the selection using a nitrocellulose filter instead of the matrix was conducted eight times (8 rounds). In the 21st round, the binding ability to the RNA pool was 22%, and the K_d value of the pool relative to the protein GST-RBD was 290 nM.

[0045] The sequences of 33 clones were determined from the RNA pool which finished the 21st round. Consequently, 24 different sequences were obtained. The sequences of the portions comprising approximately 60 bases in the 24 sequences of clones are shown in Fig. 2. They were roughly divided into two types, one having a high homology among the sequences (this group is called "group 1") and the other among which no homology, was observed (these are referred to as "group 2").

[0046] The sequences of the total sizes (approximately 100 bases) of the 8 types of RNAs (21.01 to 21.08 in Fig. 2) in group 1 are shown in sequence Nos. 1 to 8.

25 [0047] The interaction between the 10 RNAs among them and GST-RBD was examined by a binding assay using a nitrocellulose filter. The results are shown in right column of Fig. 2 in terms of RNA binding (%). As a result, the RNAs in group 1 showed the satisfactory binding to GST-RBD, whereas the RNAs in group 2 did not show the satisfactory binding. The K_d values of the RNAs shown in sequence Nos. 1 (21.01 in Fig. 2) and 7 (21.07 in Fig. 2) were both 300 nM. Meanwhile, that of the RNA in sequence No. 11 (21.11 in Fig. 2) was a micromol order (refer to Fig. 3). Fig. 3 shows percentages of binding to GST-RBD when using RNAs having sequence Nos. 1 (closed circle), 7 (closed square) and 11 (closed triangle) at various concentrations (nM).

[0048] Further, the RNA ligands were not bound to GST itself. This indicates that these RNA is bound to the RBD moiety of GST-RBD rather than to the GST moiety.

35 [0049] It was then examined whether the RNA aptamers in group 1 inhibit the interaction between Ras and RBD (refer to Fig. 4). The RNAs having sequence Nos. 1 (A in Fig. 4), 7 (B in Fig. 4), 11 (C in Fig. 4) and 12 (D in Fig. 4) (corresponding to 21.01, 21.07, 21.11 and 21.12 in Fig. 2, respectively) were tested at concentrations of 0 to 12.5 μ M. These were incubated with GST-RBD supported on a Sepharose matrix and Ras in GTP γ S or GDP. In the presence of the RNAs (lanes 3, 4 and 5 in Fig. 4; lane 3 is 20-pmol RNA, lane 4 200-pmol RNA and lane 5 2,000-pmol RNA) or in the absence of the RNAs (lanes 1 and 2 in Fig. 4; lane 1 was in the presence of GDP and lane 2 in the presence of GTP), the binding between GST-RBD and Ras was examined by immunoblotting with anti-Ras antibody RAS004.

40 [0050] In Fig. 4, "Ras" indicates Ras bound to GST-RBD, and "GST-RBD" indicates as a background that GST-RBD is solely present. As stated earlier, the RNA of sequence No. 12 (D in Fig. 4) which is scarcely bound to GST-RBD did not inhibit the binding of Ras to GST-RBD even at the concentration of 12.5 μ M. This was the same with the RNA of sequence No. 11 (C in Fig. 4) in which the K_d value was a micromol order.

45 [0051] On the other hand, the RNAs of sequence Nos. 1 (A in Fig. 4) and 7 (B in Fig. 4) in group 1 effectively inhibited the interaction between Ras and RBD. The reason is considered to be that these RNAs were bound to RBD. And, the K_d value of the GTP binding Ras and RBD of Raf-1 is 18 nM (Hermann, C., et al., J. Biol. Chem., 270, 2901 - 2905 (1995)), and the RNAs of sequence Nos. 1 and 7 have the binding ability which is 10 times lower than that. Despite this, these RNAs inhibit the interaction between Ras and Raf-1.

50 [0052] Since these RNAs do not have an affinity for Ras or a Sepharose matrix, there is no possibility that these RNAs are bound to Ras or the Sepharose matrix to inhibit the binding between GST-RBD and Ras on the matrix. This fact proved the specific binding of these RNA aptamers to RBD.

55 [0053] Another in vitro selection was conducted by using a double-stranded DNA pool obtained by synthesizing a single-stranded DNA pool (200 pmols, 1.2×10^{14} molecules) having a sequence of 5'-ggtaa tacga ctcac tatag ggagt ggagg aattc atcga ggcac-3' at the 5'-terminus and 5'-catat gcctt agcga cagca agctt ctgc-3' at the 3'-terminus and containing random 45 bases in the middle thereof and converting this single-stranded DNA pool to the double-stranded DNA pool by PCR.

[0054] Consequently, novel RNA aptamers to be bound to Raf-1 RBD could be obtained. These sequences are as follows, and shown in sequence Nos. 25 to 28 of Sequence Listing.

Sequence 25

5 gggaguggag gaaaucaucg aggcuaugu cgacuccguc uuccuucaaa ccaguuauaa 60
 auugguuuuu gcauugccu uagcgacagc aagcuucugc 100

10

Sequence 26

 gggaguggag gaaaucaucg aggcuaugacc ucccugggca guagggguua aaauuauuu 60
 15 ccuacacuuc ucaugccuua gcgacagcaa gcuucugc 98

20

Sequence 27

 gggaguggag gaaaucaucg aggcuaugu cgacuccguc uuccuucaaa ccaguuauaa 60
 auugguuuuu gcauugccu uagcgacagc 90
 25

Sequence 28

 gggaguggag gaaaucaucg aggcuaugu cgacuccguc uuccuucaaa ccaguuauaa 60
 30 auugguuuuu gcauugccu 80

35

[0055] The K_d values of the RNAs shown in sequence Nos. 25, 26 and 28 among these RNAs and GST-RBD were as follows.

RNA of sequence 25: 124 nM

RNA of sequence 26: 295 nM

RNA of sequence 28: 176 nM

40

[0056] And, the RNAs of sequences 25 to 28 all inhibited the binding between Ras and Raf-1 RBD depending on the amounts.

[0057] Sequence Nos. 25, 27 and 28 are different in the size of the 3'-terminus. From this fact, it is presumed that the RNAs of 99 to 81 bases (90 bases correspond to sequence 27) up to sequence 28 through the decrease by each one base from the 3'-side of sequence 25 also have the activity.

45

[0058] Sequences of 45 bases corresponding to the random region were shown in sequence Nos. 53 and 54.

[0059] The RNA aptamers of sequences 25 to 28 obtained here can be provided through transcription from synthetic DNAs or through synthesis.

[0060] These RNA aptamers have the stronger binding activity and are made of the smaller number of RNA bases than the above-described RNA aptamers comprising approximately 60 bases. Accordingly, these are considered to be more profitable RNA aptamers.

50

[0061] An antibody bound to Raf-1 mutant free from a kinase activity or a kinase domain of Raf-1 has been used to study the role of Ras or Raf-1 in the cell signal transduction system (Kolch, W., et al., (1991) Nature 349, 426 - 428; Kolch, W., et al., (1996) Oncogene 13, 1305 - 1314).

55

[0062] The Raf-1 mutant capable of being bound to Ras without having the kinase activity not only inhibits the Ras-dependent Raf-1 activity but also blocks the wide-ranging signal transduction systems including Ras. This is because the mutant inhibits the binding of Raf-1 and also has an influence on various downstream effectors of GTP-binding Ras.

[0063] Likewise, a monoclonal antibody bound to an epitope of a kinase domain of Raf-1 inhibits all signal transduction systems including Raf-1. This is because Raf-1 is activated not only with GTP binding Ras but also through a route having no bearing on Ras (Kolch, W., et. al., (1996) Oncogene 13, 1305 - 1314).

[0064] From this standpoint as well, it is said that the RNA aptamer to RBD in the present invention can specifically inhibit the binding between Ras and Raf-1 without having any effect on the kinase activity of Ras or Raf-1 by the other signaling pathways.

[0065] Further, the RNA aptamers of the present invention can be expressed within cells (Good, P. D., et al., (1997 Gene Ther. 4,45-54), and can be applied to a wide-ranging field.

[0066] Thus, the RNA aptamers of the present invention specifically block RBD of "target proteins of Ras", more preferably Raf-1. Not only it can be used in an agent for controlling intracellular signal transduction or a method of controlling cell signaling pathway, but also it is especially suited for the field of treatment, prevention or diagnosis of various diseases in which the signal transduction system participates.

[0067] When the nucleic acid molecular species of the present invention is used in controlling the cell signal transduction system, the nucleic acid of the present invention may directly be introduced into desired cells. It can also be introduced into cells by being inserted into viruses or the like.

[0068] Further, it is also possible that the RNA is introduced not directly but in the form of a DNA. When the nucleic acid molecular species of the present invention is used as a pharmaceutical composition, it can parenterally be administered as such, or it can be administered by being inserted into viruses or various vectors in the form of a DNA. In these administration forms, the pharmaceutical composition can also be provided using a pharmaceutically acceptable carrier.

[0069] The pharmaceutical composition of the present invention is useful for treatment, prevention or diagnosis of various diseases in which the cell signal transduction system participates, especially malignant tumors and inflammatory diseases.

Examples

[0070] The present invention is illustrated more specifically with reference to the following Examples. However, the present invention is not limited to these Examples.

Example 1 (Purification of a protein)

[0071] Glutathione S-transferase (GST) and a fusion protein of RBD (51 to 131 amino acid moiety of Raf-1) and GST (hereinafter referred to as GST-RBD) were expressed in E. coli strains BL21 and BL21DE3 respectively, and purified by chromatography using a column of glutathione-Sepharose 4B (made by Amersham Pharmacia Biotech) and HQ proth (perceptive) (Shirouzu M., et al., J. Biol. Chem., 273, 7737 - 7742 (1998)).

[0072] As wild type Ha-Ras protein, a protein obtained from E. coli strain BL21 was purified by column chromatography using DEAE-Sephacell, Sephadex G-75 and Resource QFPLC (made by Amersham Pharmacia Biotech) (Shirouzu M., et al., J. Biol. Chem., 273, 7737 - 7742 (1998); Shirouzu M., et al., Oncogene; 7, 475 - 480 (1992); Ito, Y., et al., Biochemistry, 36, 9109 - 9119 (1997)).

[0073] The purity of these proteins was identified through SDS-PAGE by staining with Coomassie Blue and/or silver. The purified protein was stored in 50% glycerol at -30°C.

Example 2 (In vitro selection)

[0074] A pool of DNAs containing random 60 bases was prepared. These DNAs have sequences 5'-GCCGGAATTCTAATACGACTCACTATAGGGAGATCAGAATAACGCTCAA-3' and 5'-TTCGACATGAGGCCCTGCAGGGCG-3' at both termini for in vitro transcription and amplification by PCR.

[0075] These RNAs were heated at 75°C for 3 minutes, then ice-cooled, and incubated in a binding buffer (5 mM MgCl₂ containing phosphate buffer physiological saline solution) containing GST-RBD and glutathione-Sepharose 4B beads. The RNAs having GST-RBD bound thereto were recovered with glutathione-Sepharose beads. The beads were cleaned with a cleaning buffer (20 mM Tris-HCl pH 7.5, 5 mM MgCl₂ and 150 mM NaCl), and the RNAs on the beads were eluted with boiling water.

[0076] The 1st through 13th rounds were conducted in this manner, provided that in order to remove the RNAs bound to GST and/or the beads alone, the RNAs were passed through glutathione-Sepharose beads having supported thereon GST only before the incubation of GST-RBD (3rd to 13th rounds) and after the elution (7th to 13th rounds).

[0077] The eluate was subjected to reverse transcription, and amplified through PCR. The RNA pool for the next round was prepared through in vitro transcription from the cDNA amplified. The 13th round and those following for selection were conducted by a filter binding method. The RNA pool (heated at 75°C for 3 minutes, and cooled to room temperature) and GST-RBD were incubated in a binding buffer at 37°C for 1 hour. The RNAs bound to GST-RBD were separated by being bound to a nitrocellulose filter. The RNAs were eluted with a buffer containing 7 M urea (Hirao, I., et al., Mol. Diversity, (under printing)). In order to remove the RNAs bound to the filter, the RNAs were passed through the filter before the amplification.

Example 3 (Nitrocellulose filter binding assay)

[0078] RNAs were obtained through the in vitro transcription with a T7 RNA polymerase using [α - 32 P]UTP. The RNA (0.8 μ M) and its protein were incubated in 50 μ l of a binding buffer at 37°C for 1 hour. A part (50 μ l) of the solution was moved on a filter, and cleaned three times with 200 μ l of a cleaning buffer.

[0079] In order to determine a dissociation constant, 1.6 nM of the RNA of which the 5' -terminus was labeled with [γ - 32 P]ATP and GST-RBD at various concentrations were incubated. The radiation dose on the filter was measured using a Fuji BAS2500 bio-imaging analyzer.

Example 4 (Inhibitory activity)

[0080] One microgram of GST-RBD in 160 μ l of a binding buffer containing 0.05% Triton X-100 was mixed with 10 μ l of a glutathione-Sepharose bead suspension in a phosphate buffer physiological saline solution. The mixture was incubated at 4°C for 30 minutes. After the gentle centrifugation, the supernatant liquid was discarded. A binding buffer solution containing 40 ng of Ras (this is bound through GTP γ S or GDP as described in a literature (Koide, H., et al., Proc. Natl. Acad. Sci. USA, 90, 8683 - 8686 (1993) and 160 μ l of the RNA was added to the remaining beads, and incubated at 4°C for 30 minutes. After the incubation, the beads were cleaned with 500 μ l of a cleaning buffer. The bound protein was eluted from the beads through separation using a Laemmli's buffer, and subjected to 15% SDS-PAGE. The product was subjected to immunoblotting with anti-Ras antibody RAS004 (Moodie, S. A., et al., Science, 260, 1658 - 1661 (1993)), and visualized with an ECL immune detector (manufactured by Amersham Pharmacia Biotech).

Example 5 (Formation of an RNA pool)

[0081] A single-stranded DNA (200 pmols, 1.2×10^{14} molecules) containing random 45 bases and having a sequence 5'-ggtaa tacga ctcac tatag ggagt ggagg aattc atcga ggcac-3' at the 5'-terminus and a sequence 5'-catat gcctt agcga cagca agctt ctgc-3' at the 3'-terminus was subjected to PCR using 2 primers, 5'-ggtaa tacga ctcac tatag ggagt ggagg aattc atcg-3' and 5'-gcaga agctt gctgt cgcta aggc-3', and then transcribed with a T7 RNA polymerase to form a first RNA pool.

Example 6 (Selection of RNA bound to Raf-1RBD)

[0082] Three micromols (1,800 pmols) of the RNA pool heated at 75°C for 3 minutes and then ice-cooled and 1 μ M (600 pmols) of GST-RBD were incubated in 600 μ l of a binding buffer at 37°C for 1 hour. The culture was filtered with a nitrocellulose filter, and the filter was cleaned three times with 300 μ l of a cleaning buffer. Thereafter, the RNAs on the filter were eluted with a buffer containing 7 M urea. After the reverse transcription, PCR was conducted at 12 cycles.

[0083] The reagents used here are as follows.

[0084] GST-RBD: A fusion protein of RBD (51 to 131 amino acid moiety of Raf-1) and glutathione-S-transferase is described in a literature (Shirouzu, M., et al., (1998) J. Biol. Chem. 273, 7737 - 7742).

binding buffer: 5 mM MgCl₂-containing phosphate buffer physiological saline solution

cleaning buffer: 20 mM Tris-HCl pH 7.5, 5 mM MgCl₂ and 150 mM NaCl

Example 7 (RNAs of sequence Nos. 25 and 26)

[0085] Selection of RNAs bound to Raf-1 RBD from the first RNA pool in Example 6, reverse transcription of RNAs to DNAs, amplification and transcription of DNAs to RNAs were repeated 10 times to obtain RNAs of sequence Nos. 25 and 26 of Sequence Listing.

Example 8 (RNAs of sequence Nos. 27 and 28)

[0086] DNAs which had a complementary sequence of an RNA of sequence No. 25 and of which the 3'-terminus side was shortened were obtained by PCR using a primer 5'-ggtaa tacga ctcac tatag ggagg ggagg aattc atcg-3' and a primer 5'-gctgt cgcta aggcata tgc taaa c-3' or 5'-aggca tatgc taaa ccaat ttata ac-3'. From these DNAs, RNAs of sequence Nos. 27 and 28 of Sequence Listing were obtained.

Example 9 (Cloning and determination of a sequence)

[0087] A DNA was cloned using a TOPO TA cloning kit, and the sequence was determined with an automated DNA sequencer.

Example 10 (Measurement of a Kd value)

5 **[0088]** An RNA (4 nM) of which the 5'-terminus was labeled and 50 to 1,250 nM of GST-RBD were incubated in 600 μ l of a binding buffer at 37°C for 30 minutes. The culture was filtered with a nitrocellulose filter, and the radioactivity on the filter was measured. The Kd value was calculated using a software: Kalleider Graph (Bell, S. D., et al., (1998) J. Biol. Chem. 273, 14309 - 14314).

Example 11 (Binding inhibition experiment)

10 **[0089]** GST-RBD (20 pmols) in 160 μ l of a binding buffer containing 0.05% Triton X-100 and 10 μ l of a phosphate buffer physiological saline solution containing glutathione-Sepharose 4B beads were mixed, and the mixture was incubated at 4°C for 30 minutes. The beads were separated, and incubated with 20 pmols of Ras and an RNA (0, 20, 100 and 200 pmols) in 160 μ l of a binding buffer (5 mM MgCl₂-containing phosphate buffer physiological saline solution) at 4°C for 30 minutes. After the beads were cleaned, the bound protein was eluted with a Laemmli's buffer, subjected to
15 SDS-PAGE, then immunoblotted using anti-Ras antibody RAS004 (Kanai, T., et al., (1987) Jpn. J. Cancer Res. 78, 1314 - 1318), and visualized with an ECL immune detector.

[0090] In the RNAs of sequence Nos. 25 to 28, the decrease in the amount of Ras was observed according to the RNA amount.

20 Industrial Applicability

[0091] The present invention provides RNAs which are specifically bound to target proteins of Ras such as Raf-1 and the like and which further inhibit the binding to Ras, and a method of specifically inhibiting an intracellular signaling pathway using these RNAs. The present invention can not only clarify the signaling pathway through the specific route
25 of cells but also provide a pharmaceutical composition having less side effects.

SEQUENCE LISTING

30 **[0092]**

<110> JAPAN SCIENCE AND TECHNOLOGY CORPORATION

<120> NUCLEIC ACID CAPABLE OF BINDING SPECIFICALLY TO RAS TARGET PROTEIN

35 <130> N79287 TAC

<140> EP 99937069.5

<141> 1999-08-13

40 <150> JP 10-242596/1998

<151> 1998-08-14

<150> JP 10-333284/1998

<151> 1998-11-24

45 <160> 61

<170> PatentIn Ver. 2.0

50 <210> 1

<211> 108

<212> RNA

<213> Artificial Sequence

<220>

55 <221> protein bind

<223> RNA aptamer

<400> 1

		gggagaucag aa metaacgcu caacugauca auggcguaca auggauucgu ucuc metaacc	60
		aaaacccua ccccuaggac ugaucgaca ugagggcccu gcagggcg	108
5		<210> 2	
		<211> 107	
		<212> RNA	
		<213> Artificial Sequence	
10		<220>	
		<221> protein bind	
		<223> RNA aptamer	
		<400> 2	
15		gggagaucag aa metaacgcu caacugauca auggcguaca auggauucgu ucuc metaacc	60
		aaaacccua ccccuaggac gaucgacau gagggcccg cagggcg	107
20		<210> 3	
		<211> 108	
		<212> RNA	
		<213> Artificial Sequence	
		<220>	
25		<221> protein bind	
		<223> RNA aptamer	
		<400> 3	
30		gggagaucag aa metaacgcu caacugauca auggcguaca auggauucgu ucuc metaacc	60
		aaaacccua ccccuaggac ugcucgaca ugagggcccu gcagggcg	108
35		<210> 4	
		<211> 108	
		<212> RNA	
		<213> Artificial Sequence	
		<220>	
		<221> protein bind	
40		<223> RNA aptamer	
		<400> 4	
45		gggagaucag aa metaacgcu caacugauca auggcguaca auggauucgc ucuc metaacc	60
		aaaacccua ccccuaggac ugcucgaca ugagggcccu gcagggcg	108
50		<210> 5	
		<211> 108	
		<212> RNA	
		<213> Artificial Sequence	
		<220>	
		<221> protein bind	
		<223> RNA aptamer	
55		<400> 5	

		gggagaucag aauaaacgcu caacugauca auggcguaca auggauucgu ucucauaacc aaaaccuuu cuccuuggac ugcucgaca ugaggcccu gcagggcg	60 108
5		<210> 6 <211> 108 <212> RNA <213> Artificial Sequence	
10		<220> <221> protein bind <223> RNA aptamer <400> 6	
15		gggagaucag aauaaacgcu caacugauca auggcguaca auggauucgu ucucauaacc aaaaccuuu cccuuggac uguuucgaca ugaggcccu gcagggcg	60 108
20		<210> 7 <211> 108 <212> RNA <213> Artificial Sequence <220>	
25		<221> protein bind <223> RNA aptamer <400> 7	
30		gggagaucag aauaaacgcu caauugacuc aauggcguac aauggauucg uucucuaaac caaaaccuu acccuugga cuguucgaca ugaggcccu gcagggcg	60 108
35		<210> 8 <211> 108 <212> RNA <213> Artificial Sequence <220>	
40		<221> protein bind <223> RNA aptamer <400> 8	
45		gggagaucag aauaaacgcu caauugaaga ucguacaug gauucgauca uaaccgaag uuuuuaaca cuuuuaccu guauucgaca ugaggcccu gcagggcg	60 108
50		<210> 9 <211> 108 <212> RNA <213> Artificial Sequence <220>	
55		<221> protein bind <223> RNA aptamer <400> 9	

	gggagaucag aaUaaacgcU caaucgaguc cacgaacauU acauauuuga acacuucagc accgaacaug cuuaguacua uccuucgaca ugaggccccu gcagggcg	60 108
5	<210> 10 <211> 108 <212> RNA <213> Artificial Sequence	
10	<220> <221> protein bind <223> RNA aptamer <400> 10	
15	gggagaucag aaUaaacgcU caauuuuacc auagccuuga gguaaacaau uuagcacacc ugaauacacg aacuaugaac ucauucgaca ugaggccccu gcagggcg	60 108
20	<210> 11 <211> 107 <212> RNA <213> Artificial Sequence <220>	
25	<221> protein bind <223> RNA aptamer <400> 11	
30	gggagaucag aaUaaacgcU caacuugagc caauuaaaag auuuacaaca agaacaugaa cgugacagcg auauauuac gauucgacau gaggccccug cagggcg	60 107
35	<210> 12 <211> 108 <212> RNA <213> Artificial Sequence <220>	
40	<221> protein bind <223> RNA aptamer <400> 12	
45	gggagaucag aaUaaacgcU caagcgacaa gcagcagaua aaguugagcg caacgccgcu acagaaccaa auuaacaugu auguucgaca ugaggccccu gcagggcg	60 108
50	<210> 13 <211> 107 <212> RNA <213> Artificial Sequence <220>	
55	<221> protein bind <223> RNA aptamer <400> 13	

	gggagaucag aaaaaacgcu caaucgaaag uaaguccgau acaacacaua accuauuuuu	60
	uagcagcgau aaacaaaaua aguucgacau gaggcccccug cagggcg	107
5	<210> 14	
	<211> 108	
	<212> RNA	
	<213> Artificial Sequence	
10	<220>	
	<221> protein bind	
	<223> RNA aptamer	
	<400> 14	
15	gggagaucag aaaaaacgcu caagcaguua uccacuugua auugaaugua gaugccauau	60
	agaguuaaua guaaucgaa uguucgaca ugaggccccc gcagggcg	108
20	<210> 15	
	<211> 108	
	<212> RNA	
	<213> Artificial Sequence	
	<220>	
25	<221> protein bind	
	<223> RNA aptamer	
	<400> 15	
30	gggagaucag aaaaaacgcu caacguagua gcacaccaug accuauuuuu ucugcuucgc	60
	aauguaccuu aacacauaau caguucgaca ugaggccccc gcagggcg	108
35	<210> 16	
	<211> 108	
	<212> RNA	
	<213> Artificial Sequence	
	<220>	
	<221> protein bind	
40	<223> RNA aptamer	
	<400> 16	
45	gggagaucag aaaaaacgcu caagaugac uaauuuuac aacagauaac cuuacucuug	60
	auaaaugcuu ugcuuuuggu uaaucgaca ugaggccccc gcagggcg	108
50	<210> 17	
	<211> 108	
	<212> RNA	
	<213> Artificial Sequence	
	<220>	
	<221> protein bind	
	<223> RNA aptamer	
55	<400> 17	

	gggagaucag aa metaacgcu caaucuucga aguccaugac ugcaaaacca gauaguccua aucucaauua ucagucccaa guauucgaca ugaggccccc gcagggcg	60 108
5	<210> 18 <211> 108 <212> RNA <213> Artificial Sequence	
10	<220> <221> protein bind <223> RNA aptamer <400> 18	
15	gggagaucag aa metaacgcu caaacacucu aaauuguggu acuaaggag uaagggaac uacgaagacg ugcaaggaua aaguucgaca ugaggccccc gcagggcg	60 108
20	<210> 19 <211> 107 <212> RNA <213> Artificial Sequence	
25	<220> <221> protein bind <223> RNA aptamer <400> 19	
30	gggagaucag aa metaacgcu caauuugccu cgacggucug cgaauagaac gcgaaccgug auuaguguac aaggauucg uuuucgacau gaggccccug cagggcg	60 107
35	<210> 20 <211> 106 <212> RNA <213> Artificial Sequence	
40	<220> <221> protein bind <223> RNA aptamer <400> 20	
45	gggagaucag aa metaacgcu caagucgcag cagaaauauc aucgaaaac cucaauugca ucucauguau aucuagucca auucgacau agggccccug agggcg	60 106
50	<210> 21 <211> 105 <212> RNA <213> Artificial Sequence	
55	<220> <221> protein bind <223> RNA aptamer <440> 21	

	gggagaucag aa metaacgcu caacgaacau cuggaguaau caucuaaau accucauuua ccuuuacacu uucuaaacua uucgacauga ggccccugca gggcg	60 105
5	<210> 22 <211> 108 <212> RNA <213> Artificial Sequence	
10	<220> <221> protein bind 423> RNA aptamer <400> 22	
15	gggagaucag aa metaacgcu caaggguuag ggugagcagu ucaagauggu aacuggcauu cauuugaaga aaggguuggua gacuucgaca ugaggcccu gcagggcg	60 108
20	<210> 23 <211> 108 <212> RNA <213> Artificial Sequence	
25	<220> <221> protein bind <223> RNA aptamer <400> 23	
30	gggagaucag aa metaacgcu caaggguuag ggugagcagu ucaagauggu aaccggcauu cauuugaaga aaggguuggua aacuucgaca ugaggcccu gcagggcg	60 108
35	<210> 24 <211> 101 <212> RNA <213> Artificial Sequence	
40	<220> <221> protein bind <223> RNA aptamer <400> 24	
45	gggagaucag aa metaacgcu caacuuggug uaguguucaa gugagauaua guuaaagguu auuguugugc gaacggguucg acaugaggcc ccugcagggc g	60 101
50	<210> 25 <211> 100 <212> RNA <213> Artificial Sequence	
55	<220> <221> protein bind <223> RNA aptamer <400> 25	

	gggaguggag gaaaucaucg aggcauauu cgacuccguc uuccuucaaa ccaguuauaa auuggguuuu gcauauccu uagcgacagc aagcuucugc	60 100
5	<210> 26 <211> 98 <212> RNA <213> Artificial Sequence	
10	<220> <221> protein bind <223> RNA aptamer <400> 26	
15	gggaguggag gaaaucaucg aggcauauu ucccugggca guagggguua aaauuauuu ccuacacuuu ucauugccuu gcgacagcaa gcuucugc	60 98
20	<210> 27 <211> 90 <212> RNA <213> Artificial Sequence <220>	
25	<221> protein bind <223> RNA aptamer <400> 27	
30	gggaguggag gaaaucaucg aggcauauu cgacuccguc uuccuucaaa ccaguuauaa auuggguuuu gcauauccu uagcgacagc	60 90
35	<210> 28 <211> 80 <212> RNA <213> Artificial Sequence <220>	
40	<221> protein bind <223> RNA aptamer <400> 28	
45	gggaguggag gaaaucaucg aggcauauu cgacuccguc uuccuucaaa ccaguuauaa auuggguuuu gcauauccu	60 80
50	<210> 29 <211> 60 <212> RNA <213> Artificial Sequence <220>	
55	<221> protein bind <223> RNA aptamer <400> 29	

	cugaucaaug gcguacaaug gauucguucu cauaaccaa acccuuaccc cuuggacuga	60
5	<210> 30 <211> 59 <212> RNA <223> Artificial Sequence <220>	
10	<221> protein bind <223> RNA aptamer <400> 30	
15	cugaucaaug gcguacaaug gauucguucu cauaaccaa acccuuaccc cuuggacuga	59
20	<210> 31 <211> 60 <212> RNA <213> Artificial Sequence <220> <221> protein bind <223> RNA aptamer	
25	<400> 31	
30	cugaucaaug gcguacaaug gauucguucu cauaaccaa acccuuaccc cuuggacugc	60
35	<210> 32 <211> 60 <212> RNA <213> Artificial Sequence <220> <221> protein bind <223> RNA aptamer	
40	<400> 32	
45	cugaucaaug gcguacaaug gauucgcucu cauaaccaa acccuuaccc cuuggacugc	60
50	<210> 33 <211> 60 <212> RNA <213> Artificial Sequence <220> <221> protein bind <223> RNA aptamer	
55	<400> 33	
	cugaucaaug gcguacaaug gauucguucu cauaaccaa acccuuacuc cuuggacugc	60
	<210> 34 <211> 60 <212> RNA	

EP 1 026 243 B1

<213> Artificial Sequence
 <220>
 <221> protein bind
 <223> RNA aptamer
 5 <400> 34

cugaucaaug gcguacaug gauucguucu cauaaccaa acccuaccc cuuggacugu 60

10
 <210> 35
 <211> 60
 <212> RNA
 <213> Artificial Sequence
 15 <220>
 <221> protein bind
 <223> RNA aptamer
 <400> 35

20
 uugacucaau ggcguacaau ggauucguuc ucauaaccaa aaccuuacc ccuuggacug 60

25
 <210> 36
 <211> 60
 <212> RNA
 <213> Artificial Sequence
 <220>
 <221> protein bind
 30 <223> RNA aptamer
 <400> 36

35
 uugaagaucg uacauggau ucgaucuaa cccgaaguuu uuaaacacuc uuuaaccugua 60

40
 <210> 37
 <211> 60
 <212> RNA
 <213> Artificial Sequence
 <220>
 <221> protein bind
 <223> RNA aptamer
 45 <400> 37

ucgaguccac gaacauuaca uauuugaaca cuucagcacc gaacaugcuu aguacuaucc 60

50
 <210> 38
 <211> 60
 <212> RNA
 <213> Artificial Sequence
 <220>
 55 <221> protein bind
 <223> RNA aptamer
 <400> 38

	uauuaccaua gccuugaggu aaacaaauua gcacaccuga auacacgaac uaugaacuca	60
5	<210> 39 <211> 59 <212> RNA <213> Artificial Sequence <220>	
10	<221> protein bind <223> RNA aptamer <400> 39	
15	cuugagccaa uaaaaagauu uacaacaaga acaugaacgu gacagcgaua auaaauacga	59
20	<210> 40 <211> 60 <212> RNA <213> Artificial Sequence <220> <221> protein bind <223> RNA aptamer	
25	<400> 40	
	gcgacaagca gcagauaaag uugagcgcaa cgccgcuaca gaaccaaaau aacauguang	60
30	<210> 41 <211> 59 <212> RNA <213> Artificial Sequence <220>	
35	<221> protein bind <223> RNA aptamer <400> 41	
40	ucgaaaguua guccgauaca acacauaacc uauuuuuuag cagcgauau acaauaag	59
45	<210> 42 <211> 60 <212> RNA <213> Artificial Sequence <220> <221> protein bind <223> RNA aptamer	
50	<400> 42	
	gcaguaaucc acuuguaauu gaauaguagau gccauauaga guuuuuagua auccgaauug	60
55	<210> 43 <211> 60 <212> RNA	

<213> Artificial Sequence
 <220>
 <221> protein bind
 <223> RNA aptamer
 5 <400> 43

cguaguagca caccaugacc uauuaaauc ugcucgcaau guaccuuuac acauaaucag 60

10

<210> 44
 <211> 60
 <212> RNA
 <213> Artificial Sequence
 15 <220>
 <221> protein bind
 <223> RNA aptamer
 <400> 44

20

gaaugacuaa uaaauacaac agauaaccuu acucuugaua aaugcuuugc uuugguuaa 60

25

<210> 45
 <211> 60
 <212> RNA
 <213> Artificial Sequence
 <220>
 <221> protein bind
 30 <223> RNA aptamer
 <400> 45

35

ucucgaagu ccaugacugc aaaaccagau aguccuauc ucauuuauca gucccaagua 60

40

<210> 46
 <211> 60
 <212> RNA
 <213> Artificial Sequence
 <220>
 <221> protein bind
 <223> RNA aptamer
 45 <400> 46

50

acacucuaaa uuugguacu aaggagaua gggcaacuac gaagacgugc aaggauaaag 60

55

<210> 47
 <211> 59
 <212> RNA
 <213> Artificial Sequence
 <220>
 <221> protein bind
 <223> RNA aptamer
 <400> 47

		uuugccucga cggucugcga auagaacgcg aaccgugauu aguguacaag gauucgguu	59
5	<210> 48 <211> 58 <212> RNA <213> Artificial Sequence <220>		
10	<221> protein bind <223> RNA aptamer <400> 48		
15		gucgcagcag aaauaucac gcaaaaccuc aaugcaucu cauguauauc uaguccaa	58
20	<210> 49 <211> 57 <212> RNA <213> Artificial Sequence <220>		
25	<221> protein bind <223> RNA aptamer <400> 49		
30		cgaacaucug gaguaaucau cuuaauaacc ucauuaccu uuacacuuuc uaaacua	57
35	<210> 50 <211> 60 <212> RNA <213> Artificial Sequence <220>		
40	<221> protein bind <223> RNA aptamer <400> 50		
45		ggguaagggg gagcaguuca agaugguaac uggcauucan uugaagaaag guugguagac	60
50	<210> 51 <211> 60 <212> RNA <213> Artificial Sequence <220>		
55	<221> protein bind <223> RNA aptamer <400> 51		
		ggguaagggg gagcaguuca agaugguaac cggcauucan uugaagaaag guugguaaac	60
	<210> 52 <211> 53 <212> RNA		

	<213> Artificial Sequence	
	<220>	
	<221> protein bind	
	<223> RNA aptamer	
5	<400> 52	
	cuugguguag uguucaagug agauauagua uaagguuuuu guugugcgaa cgg	53
10	<210> 53	
	<211> 45	
	<212> RNA	
	<213> Artificial Sequence	
15	<220>	
	<221> protein bind	
	<223> RNA aptamer	
	<400> 53	
20	augucgacuc cgucuuccuu caaaccaguu auaaaauuggu uuuag	45
	<210> 54	
	<211> 45	
25	<212> RNA	
	<213> Artificial Sequence	
	<220>	
	<221> protein bind	
	<223> RNA aptamer	
30	<400> 54	
	gaccucccggu ggcaguaggg guaaaaauua ucuuccuaca cuucu	45
	--- --	
35	<210> 55	
	<211> 23	
	<212> RNA	
	<213> Artificial Sequence	
	<220>	
40	<221> prim transcript	
	<223> primer for cDNA	
	<400> 55	
45	gggagaucag auaaaacgcu caa	23
	<210> 56	
	<211> 25	
50	<212> RNA	
	<213> Artificial Sequence	
	<220>	
	<221> prim transcript	
	<223> primer for cDNA	
55	<400> 56	

	uucgacauga ggccccugca gggcg	25
5	<210> 57 <211> 50 <212> DNA <213> Artificial Sequence <220>	
10	<221> primer bind <223> PCR primer <400> 57	
15	ggcgggaattc taatacgact cactataggg agatcagaat aaacgctcaa	50
20	<210> 58 <211> 25 <212> DNA <213> Artificial Sequence <220>	
25	<221> primer bind <223> PCR primer <400> 58	
30	cgccctgcag gggcctcatg tcgaa _ _ _ _ _	25
35	<210> 59 <211> 55 <212> DNA <213> Artificial Sequence <220>	
40	<221> primer bind <223> PCR primer <400> 59	
45	ggtaatacga ctactatag ggagtggagg aattcatcga ggcatt	45
50	<210> 60 <211> 29 <212> DNA <213> Artificial Sequence <220>	
55	<221> primer bind <223> PCR primer <400> 60	
	catatgcctt agcgacagca agcttctgc	29
	<210> 61 <211> 108	

<212> RNA
 <213> Artificial Sequence
 <220>
 <221> protein bind
 <223> RNA aptamer
 <400> 61

5

10

gggagaucag aauaaacgcu caannnnnnnn nnnnnnnnnnn nnnnnnnnnnn nnnnnnnnnnn 60
 nnnnnnnnnnn nnnnnnnnnnn nnnuucgaca ugaggccccc gcagggcg 108

15

Claims

20

1. An RNA capable of being specifically bound to a Ras binding domain of Raf-1, wherein the RNA contains at least one sequence selected from sequence Nos. 1 to 8 or 25 to 28 of the Sequence Listing or a sequence of 81 to 99 bases obtainable by deletion of a base or bases from the 3'-terminus of sequence No. 25.

25

2. A nucleic acid having a complementary base sequence to the RNA defined in claim 1.

3. An agent for controlling cell signal transduction which agent is made of the RNA according to claim 1 or the nucleic acid according to claim 2.

25

4. A method of controlling cell signal transduction *in vitro* using the RNA according to claim 1 or a nucleic acid according to claim 2.

30

5. A pharmaceutical composition containing the RNA according to claim 1 or a nucleic acid according to claim 2.

6. Use of a pharmaceutical composition according to claim 5 in the manufacture of a medicament for use in the treatment of cancer or an inflammatory disease.

35

Revendications

40

1. ARN pouvant être lié d'une manière spécifique à un domaine de liaison Ras de Raf-1, l'ARN contenant au moins une séquence choisie parmi les séquences N° 1 à 8 ou 25 à 28 de la Liste des Séquences ou une séquence de 81 à 99 bases pouvant être obtenue par délétion d'une ou plusieurs bases à partir du site 3'-terminal de la séquence n° 25.

45

2. Acide nucléique ayant une séquence de bases complémentaire de l'ARN défini dans la revendication 1.

3. Agent destiné à réguler la transduction des signaux cellulaires, lequel agent est constitué de l'ARN selon la revendication 1 ou de l'acide nucléique selon la revendication 2.

50

4. Procédé de régulation de la transduction des signaux cellulaires *in vitro* par utilisation de l'ARN selon la revendication 1 ou d'un acide nucléique selon la revendication 2.

5. Composition pharmaceutique contenant l'ARN selon la revendication 1 ou un acide nucléique selon la revendication 2

6. Utilisation d'une composition pharmaceutique selon la revendication 5 pour préparer un médicament destiné à être utilisé dans le traitement du cancer ou d'une maladie inflammatoire.

55

Patentansprüche

1. RNA, die fähig ist, durch eine Ras-Bindungsdomäne von Raf-1 spezifisch gebunden zu werden, wobei die RNA mindestens eine Sequenz enthält, ausgewählt aus Sequenz Nrn. 1 bis 8 oder 25 bis 28 der Sequenzliste oder eine

EP 1 026 243 B1

Sequenz von 81 bis 99 Basen, die durch Deletion einer Base oder von Basen vom 3'-Terminus der Sequenz Nr. 25 erhaltbar ist.

- 5
2. Nukleinsäure mit einer komplementären Basensequenz zu der in Anspruch 1 definierten RNA.
3. Agens für die Kontrolle der Zellsignaltransduktion, welches Agens aus der RNA nach Anspruch 1 oder der Nukleinsäure nach Anspruch 2 hergestellt ist.
- 10
4. Methode zur Kontrolle der Zellsignaltransduktion *in vitro* unter Verwendung der RNA nach Anspruch 1 oder einer Nukleinsäure nach Anspruch 2.
5. Pharmazeutische Zusammensetzung, enthaltend die RNA nach Anspruch 1 oder eine Nukleinsäure nach Anspruch 2.
- 15
6. Verwendung einer pharmazeutischen Zusammensetzung nach Anspruch 5 in der Herstellung eines Medikaments zur Anwendung in der Behandlung von Krebs oder einer entzündlichen Erkrankung.

20

25

30

35

40

45

50

55

Starting RNA:
5' -GGGAGAUCAAGAAUAACGCCUAA[-60N-]UUCGACAUGAGGCCCCUUGCAGGGCG-3'

PCR primer 1:
5' -GCCGGAATTCAATACGACTCACTATAGGAGATCAGAAATAACGCTCA-3',
EcoRI T7 promoter

PCR primer 2:
5' -CGCCCTGCAGGGGCTCATGTGAA-3',
PstI

Figure 1

RNA clone ^a	Sequence ^b	RNA Bound (%) ^c
Group 1		
21.01 (6)	CUGAUCAAUGGGGUAACAAGGAUUCGUUUCUCAUAACCAAAACCCUUAACCCUUGGACUGA	42
21.02	CUGAUCAAUGGGGUAACAAGGAUUCGUUUCUCAUAACCAAAACCCUUAACCCUUGGACUGA	
21.03 (4)	CUGAUCAAUGGGGUAACAAGGAUUCGUUUCUCAUAACCAAAACCCUUAACCCUUGGACUGC	
21.04	CUGAUCAAUGGGGUAACAAGGAUUCGUUUCUCAUAACCAAAACCCUUAACCCUUGGACUGC	46
21.05	CUGAUCAAUGGGGUAACAAGGAUUCGUUUCUCAUAACCAAAACCCUUAACCCUUGGACUGC	
21.06	CUGAUCAAUGGGGUAACAAGGAUUCGUUUCUCAUAACCAAAACCCUUAACCCUUGGACUGC	46
21.07	UUGACUCAUUGGGGUAACAAGGAUUCGUUUCUCAUAACCAAAACCCUUAACCCUUGGACUG	46
21.08	UUGAAGAUCGUAACAAGGAUUCG--AUCAUAAACCCGAAGUUUUUAAACACUCUUUACUGUA	12
Group 2		
21.09	UUGAGUCCACGAACAUAUAUUUUGAACACUUCAGCACCGAACCAUGCUUAGUAUAUCC	
21.10	UAUUAACAUAAGCCUUGAGGUAACAUAUUUAGCACACCUUGAAUACACGAAUAUGAACUCA	3
21.11 (2)	CUUGAGCCAAUUAAGAUAUUUACAACAAGAACAUAGACGUGACAGCGAUAUAUAUACGA	0.2
21.12	GCGACAGCAGCAGAUAAAGUUGAGCGCAACGCCGCUACAGAACCAAAUUAACAUGUAUG	
21.13	UCGAAAGUAAGUCCGAUACAACAUAACCUAUUAUUUAGCAGCGAUAUAACAUAUAG	
21.14	GCAGUAUCCACUUGUAUUUGAAUGUAUGGCAUAUAAGAUUAUAUAUAUUAUCCGAAUUG	
21.15	CGUAGUAGCACACCAUGACCUUAUAUUCUGCUUCGCAUUGUACCUUAACACAUUAUCAG	1
21.16	GAAUGACUAUAUAUAACAAGUAACCUUAUCUUAUAUAUAUAUAUAUAUAUAUAUAUAUA	
21.17	UCUUCGAAGUCCAUGACUGCAAAACCAAGUAUAGUCCUAUUCUAAUAUAUACAGUCCCAAGUA	0.5
21.18	ACACUCUAAAUUGUGGUACUAAGGAGUAAGGGCAACUACGAAGACGUGCAAGGAUAUAG	
21.19	UUUGCCUCCGACGGUCUGCGAAUAAGAACCGGAAACCGUGAUUAUAUAUAUAUAUAUAUAUA	0.7
21.20	GUCCGACGAGAAUA	
21.21	CGAACAUUCUGGAGUA	1.3
21.22	GGGUAAGGGUGAGCAGUUA	
21.23	GGGUAAGGGUGAGCAGUUA	
21.24	CUUGGUGUAGUGUUA	

Figure 2




Figure 3




Figure 4

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☐ BLACK BORDERS

☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES

☒ FADED TEXT OR DRAWING

☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING

☐ SKEWED/SLANTED IMAGES

☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS

☐ GRAY SCALE DOCUMENTS

☐ LINES OR MARKS ON ORIGINAL DOCUMENT

☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY

☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.